Simon Whitaker MA VetMB MRCVS
Paul McCarthy BVSc MANZCVSc (Med)
Jonathon Nott BVSc (Hons) MANZCVSc (Surg)
Sue Graykowski BVSc
Sacha Woodburn BVSc (Hons)
Tiarna Barclay BVSc
Kimberley Chainey BVSc
Melanie Ngo BvetBiol/DVM
Clare Bumak BVSc (Hons1)



Adamstown NSW 2289
Telephone 02 4957 2269
Facsimile 02 4952 8666
Emailbrunkerroadvets@bigpond.com.au

Website www.brunkerroadvets.com.au

278 Brunker Road

Post Operative Instructions for Dental Procedure

- Your pet has had a dental procedure that included a general anaesthetic.
- After an anaesthetic, animals may be unsteady on their feet, and may shake or tremble. Keep them warm and safely confined for 24 hours after the procedure.
- Provide fresh water. Soft food can be offered.
- Hair may have been shaved from a foreleg for blood collection and/or administration of intravenous anaesthetic.
- It is normal for animals to have a cough for 1 or 2 days. This is due to a tube that was passed into their airway to deliver oxygen & prevent fluid entering the lungs.
- The procedure has included ultrasonic scaling to clean tartar away and possibly extractions to remove any loose or infected teeth.
- For the next day or so you may find your pet will:
 - Salivate with a small amount of blood.
 - Lick lips or paw at mouth.
 - Be tender to touch around the mouth and head.

These signs should subside after a couple of days.

- We advise soft food to be given for 3-4 days after a dental, then back to their normal diet. Owners often worry that their pets will have trouble eating if multiple teeth have been extracted but the gums heal well and form a hard surface that functions well for chewing food.
- You may be sent home with medications in tablet or liquid form. READ THE DIRECTIONS
 CAREFULLY. Please let us know if you have difficulty in giving your pet the medication prescribed.
- If bleeding persists, or your pet seems depressed or refuses to eat for more than 2 days, please give us a call.

Prevention

- Tartar forms on the teeth for many reasons :- individual mouth chemistry, genetics, infections and a lack of chewy food in the diet can all contribute.
- Some dry foods are formulated to reduce tartar development (eg Hill's T/D).
- Sometimes brushing teeth with fingerstall brushes or pet toothbrushes, pet toothpaste or the use of teeth clean additives to their water
- There are also rawhide chews and purpose teeth cleaning chews available such as Greenies.

If you have problems please contact the Brunker Road Veterinary Centre on 49572269

